

40 Minutes—Graphing Calculator Required

Notes: (1) The exact numerical value of the correct answer does not always appear among the choices given. When this happens, select from among the choices the number that best approximates the exact numerical value.

(2) Unless otherwise specified, the domain of a function  $f$  is assumed to be the set of all real numbers  $x$  for which  $f(x)$  is a real number.

76. Which of the following sequences converge?

I.  $\left\{ \frac{5n}{2n-1} \right\}$

II.  $\left\{ \frac{e^n}{n} \right\}$

III.  $\left\{ \frac{e^n}{1+e^n} \right\}$

- (A) I only    (B) II only    (C) I and II only    (D) I and III only    (E) I, II, and III

77. When the region enclosed by the graphs of  $y = x$  and  $y = 4x - x^2$  is revolved about the  $y$ -axis, the volume of the solid generated is given by

(A)  $\pi \int_0^3 (x^3 - 3x^2) dx$

(B)  $\pi \int_0^3 (x^2 - (4x - x^2)^2) dx$

(C)  $\pi \int_0^3 (3x - x^2)^2 dx$

(D)  $2\pi \int_0^3 (x^3 - 3x^2) dx$

(E)  $2\pi \int_0^3 (3x^2 - x^3) dx$

78.  $\lim_{h \rightarrow 0} \frac{\ln(e+h)-1}{h}$  is

- (A)  $f'(e)$ , where  $f(x) = \ln x$   
 (B)  $f'(e)$ , where  $f(x) = \frac{\ln x}{x}$   
 (C)  $f'(1)$ , where  $f(x) = \ln x$   
 (D)  $f'(1)$ , where  $f(x) = \ln(x+e)$   
 (E)  $f'(0)$ , where  $f(x) = \ln x$

79. The position of an object attached to a spring is given by  $y(t) = \frac{1}{6} \cos(5t) - \frac{1}{4} \sin(5t)$ , where  $t$  is time in seconds. In the first 4 seconds, how many times is the velocity of the object equal to 0?

- (A) Zero  
 (B) Three  
 (C) Five  
 (D) Six  
 (E) Seven

80. Let  $f$  be the function given by  $f(x) = \cos(2x) + \ln(3x)$ . What is the least value of  $x$  at which the graph of  $f$  changes concavity?

- (A) 0.56            (B) 0.93            (C) 1.18            (D) 2.38            (E) 2.44

81. Let  $f$  be a continuous function on the closed interval  $[-3, 6]$ . If  $f(-3) = -1$  and  $f(6) = 3$ , then the Intermediate Value Theorem guarantees that

- (A)  $f(0) = 0$   
 (B)  $f'(c) = \frac{4}{9}$  for at least one  $c$  between  $-3$  and  $6$   
 (C)  $-1 \leq f(x) \leq 3$  for all  $x$  between  $-3$  and  $6$   
 (D)  $f(c) = 1$  for at least one  $c$  between  $-3$  and  $6$   
 (E)  $f(c) = 0$  for at least one  $c$  between  $-1$  and  $3$

1997 AP Calculus BC:  
Section I, Part B

82. If  $0 \leq x \leq 4$ , of the following, which is the greatest value of  $x$  such that  $\int_0^x (t^2 - 2t) dt \geq \int_2^x t dt$ ?

- (A) 1.35      (B) 1.38      (C) 1.41      (D) 1.48      (E) 1.59

83. If  $\frac{dy}{dx} = (1 + \ln x)y$  and if  $y = 1$  when  $x = 1$ , then  $y =$

- (A)  $e^{\frac{x^2-1}{x^2}}$   
(B)  $1 + \ln x$   
(C)  $\ln x$   
(D)  $e^{2x+x \ln x - 2}$   
(E)  $e^{x \ln x}$

84.  $\int x^2 \sin x dx =$

- (A)  $-x^2 \cos x - 2x \sin x - 2 \cos x + C$   
(B)  $-x^2 \cos x + 2x \sin x - 2 \cos x + C$   
(C)  $-x^2 \cos x + 2x \sin x + 2 \cos x + C$   
(D)  $-\frac{x^3}{3} \cos x + C$   
(E)  $2x \cos x + C$

85. Let  $f$  be a twice differentiable function such that  $f(1) = 2$  and  $f(3) = 7$ . Which of the following must be true for the function  $f$  on the interval  $1 \leq x \leq 3$ ?

- I. The average rate of change of  $f$  is  $\frac{5}{2}$ .  
II. The average value of  $f$  is  $\frac{9}{2}$ .  
III. The average value of  $f'$  is  $\frac{5}{2}$ .
- (A) None  
(B) I only  
(C) III only  
(D) I and III only  
(E) II and III only

86. 
$$\int \frac{dx}{(x-1)(x+3)} =$$

(A)  $\frac{1}{4} \ln \left| \frac{x-1}{x+3} \right| + C$

(B)  $\frac{1}{4} \ln \left| \frac{x+3}{x-1} \right| + C$

(C)  $\frac{1}{2} \ln |(x-1)(x+3)| + C$

(D)  $\frac{1}{2} \ln \left| \frac{2x+2}{(x-1)(x+3)} \right| + C$

(E)  $\ln |(x-1)(x+3)| + C$

---

87. The base of a solid is the region in the first quadrant enclosed by the graph of  $y = 2 - x^2$  and the coordinate axes. If every cross section of the solid perpendicular to the  $y$ -axis is a square, the volume of the solid is given by

(A)  $\pi \int_0^2 (2-y)^2 dy$

(B)  $\int_0^2 (2-y) dy$

(C)  $\pi \int_0^{\sqrt{2}} (2-x^2)^2 dx$

(D)  $\int_0^{\sqrt{2}} (2-x^2)^2 dx$

(E)  $\int_0^{\sqrt{2}} (2-x^2) dx$

88. Let  $f(x) = \int_0^{x^2} \sin t \, dt$ . At how many points in the closed interval  $[0, \sqrt{\pi}]$  does the instantaneous rate of change of  $f$  equal the average rate of change of  $f$  on that interval?

- (A) Zero
- (B) One
- (C) Two
- (D) Three
- (E) Four

89. If  $f$  is the antiderivative of  $\frac{x^2}{1+x^5}$  such that  $f(1) = 0$ , then  $f(4) =$

- (A)  $-0.012$       (B)  $0$       (C)  $0.016$       (D)  $0.376$       (E)  $0.629$

90. A force of 10 pounds is required to stretch a spring 4 inches beyond its natural length. Assuming Hooke's law applies, how much work is done in stretching the spring from its natural length to 6 inches beyond its natural length?

- (A) 60.0 inch-pounds
- (B) 45.0 inch-pounds
- (C) 40.0 inch-pounds
- (D) 15.0 inch-pounds
- (E) 7.2 inch-pounds