

Section 3-1 : Parametric Equations and Curves

1. Eliminate the parameter for the following set of parametric equations, sketch the graph of the parametric curve and give any limits that might exist on x and y .

$$x = 4 - 2t \quad y = 3 + 6t - 4t^2$$

Step 1

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we can do that by solving the x equation for t and plugging that into the y equation.

Doing that gives (we'll leave it to you to verify all the algebra bits...),

$$t = \frac{1}{2}(4 - x) \quad \rightarrow \quad y = 3 + 6\left[\frac{1}{2}(4 - x)\right] - 4\left[\frac{1}{2}(4 - x)\right]^2 = -x^2 + 5x - 1$$

Step 2

Okay, from this it looks like we have a parabola that opens downward. To sketch the graph of this we'll need the x -intercepts, y -intercept and most importantly the vertex.

For notational purposes let's define $f(x) = -x^2 + 5x - 1$.

The x -intercepts are then found by solving $f(x) = 0$. Doing this gives,

$$-x^2 + 5x - 1 = 0 \quad \rightarrow \quad x = \frac{-5 \pm \sqrt{(5)^2 - 4(-1)(-1)}}{2(-1)} = \frac{5 \pm \sqrt{21}}{2} = 0.2087, 4.7913$$

The y -intercept is : $(0, f(0)) = (0, -1)$.

Finally, the vertex is,

$$\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right)\right) = \left(\frac{-5}{2(-1)}, f\left(\frac{5}{2}\right)\right) = \left(\frac{5}{2}, \frac{21}{4}\right)$$

Step 3

Before we sketch the graph of the parametric curve recall that all parametric curves have a direction of motion, *i.e.* the direction indicating increasing values of the parameter, t in this case.

There are several ways to get the direction of motion for the curve. One is to plug in values of t into the parametric equations to get some points that we can use to identify the direction of motion.

Here is a table of values for this set of parametric equations.

t	x	y
-1	6	-7
0	4	3
$\frac{3}{4}$	$\frac{5}{2}$	$\frac{21}{4}$
1	2	5
2	0	-1
3	-2	-15

Note that $t = \frac{3}{4}$ is the value of t that give the vertex of the parabola and is not an obvious value of t to use! In fact, this is a good example of why just using values of t to sketch the graph is such a bad way of getting the sketch of a parametric curve. It is often very difficult to determine a good set of t 's to use.

For this table we first found the vertex t by using the fact that we actually knew the coordinates of the vertex (the x -coordinate for this example was the important one) as follows,

$$x = \frac{5}{2} : \frac{5}{2} = 4 - 2t \quad \rightarrow \quad t = \frac{3}{4}$$

Once this value of t was found we chose several values of t to either side for a good representation of t for our sketch.

Note that, for this case, we used the x -coordinates to find the value of the t that corresponds to the vertex because this equation was a linear equation and there would be only one solution for t . Had we used the y -coordinate we would have had to solve a quadratic (not hard to do of course) that would have resulted in two t 's. The problem is that only one t gives the vertex for this problem and so we'd need to then check them in the x equation to determine the correct one. So, in this case we might as well just go with the x equation from the start.

Also note that there is an easier way (probably – it will depend on you of course) to determine direction of motion. Take a quick look at the x equation.

$$x = 4 - 2t$$

Because of the minus sign in front of the t we can see that as t increases x must decrease (we can verify with a quick derivative/Calculus I analysis if we want to). This means that the graph must be tracing out from right to left as the table of values above in the table also indicates.

Using a quick Calculus analysis of one, or both, of the parametric equations is often a better and easier method for determining the direction of motion for a parametric curve. For “simple” parametric equations we can often get the direction based on a quick glance at the parametric equations and it avoids having to pick “nice” values of t for a table.

Step 4

We could sketch the graph at this point, but let's first get any limits on x and y that might exist.

Because we have a parabola that opens downward and we've not restricted t 's in any way we know that we'll get the whole parabola. This in turn means that we won't have any limits at all on x but y must satisfy $y \leq \frac{21}{4}$ (remember the y -coordinate of the vertex?).


So, formally here are the limits on x and y .

$$-\infty < x < \infty \qquad y \leq \frac{21}{4}$$

Note that having the limits on x and y will often help with the actual graphing step so it's often best to get them prior to sketching the graph. In this case they don't really help as we can sketch the graph of a parabola without these limits, but it's just good habit to be in so we did them first anyway.

Step 5

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included the points from our table because we had them but we won't always include them as we are often only interested in the sketch itself and the direction of motion.

2. Eliminate the parameter for the following set of parametric equations, sketch the graph of the parametric curve and give any limits that might exist on x and y .

$$x = 4 - 2t \qquad y = 3 + 6t - 4t^2 \qquad 0 \leq t \leq 3$$

Step 1

Before we get started on this problem we should acknowledge that this problem is really just a restriction on the first problem (*i.e.* it is the same problem except we restricted the values of t to use). As such we could just go back to the first problem and modify the sketch to match the restricted values of t to get a quick solution and in general that is how a problem like this would work.

However, we're going to approach this solution as if this was its own problem because we won't always have the more general problem worked ahead of time. So, let's proceed with the problem assuming we haven't worked the first problem in this section.

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we can do that by solving the x equation for t and plugging that into the y equation.

Doing that gives (we'll leave it to you to verify all the algebra bits...),

$$t = \frac{1}{2}(4-x) \quad \rightarrow \quad y = 3 + 6\left[\frac{1}{2}(4-x)\right] - 4\left[\frac{1}{2}(4-x)\right]^2 = -x^2 + 5x - 1$$

Step 2

Okay, from this it looks like we have a parabola that opens downward. To sketch the graph of this we'll need the x -intercepts, y -intercept and most importantly the vertex.

For notational purposes let's define $f(x) = -x^2 + 5x - 1$.

The x -intercepts are then found by solving $f(x) = 0$. Doing this gives,

$$-x^2 + 5x - 1 = 0 \quad \rightarrow \quad x = \frac{-5 \pm \sqrt{(5)^2 - 4(-1)(-1)}}{2(-1)} = \frac{5 \pm \sqrt{21}}{2} = 0.2087, 4.7913$$

The y -intercept is : $(0, f(0)) = (0, -1)$.

Finally, the vertex is,

$$\left(-\frac{b}{2a}, f\left(-\frac{b}{2a}\right)\right) = \left(\frac{-5}{2(-1)}, f\left(\frac{5}{2}\right)\right) = \left(\frac{5}{2}, \frac{21}{4}\right)$$

Step 3

Before we sketch the graph of the parametric curve recall that all parametric curves have a direction of motion, *i.e.* the direction indicating increasing values of the parameter, t in this case.

There are several ways to get the direction of motion for the curve. One is to plug in values of t into the parametric equations to get some points that we can use to identify the direction of motion.

Here is a table of values for this set of parametric equations. Also note that because we've restricted the value of t for this problem we need to keep that in mind as we chose values of t to use.

t	x	y
0	4	3
$\frac{3}{4}$	$\frac{5}{2}$	$\frac{21}{4}$
1	2	5
2	0	-1
3	-2	-15

Note that $t = \frac{3}{4}$ is the value of t that give the vertex of the parabola and is not an obvious value of t to use! In fact, this is a good example of why just using values of t to sketch the graph is such a bad way of getting the sketch of a parametric curve. It is often very difficult to determine a good set of t 's to use.

For this table we first found the vertex t by using the fact that we actually knew the coordinates of the vertex (the x -coordinate for this example was the important one) as follows,

$$x = \frac{5}{2} : \frac{5}{2} = 4 - 2t \quad \rightarrow \quad t = \frac{3}{4}$$

Once this value of t was found we chose several values of t to either side for a good representation of t for our sketch.

Note that, for this case, we used the x -coordinates to find the value of the t that corresponds to the vertex because this equation was a linear equation and there would be only one solution for t . Had we used the y -coordinate we would have had to solve a quadratic (not hard to do of course) that would have resulted in two t 's. The problem is that only one t gives the vertex for this problem and so we'd need to then check them in the x equation to determine the correct one. So, in this case we might as well just go with the x equation from the start.

Also note that there is an easier way (probably – it will depend on you of course) to determine direction of motion. Take a quick look at the x equation.

$$x = 4 - 2t$$

Because of the minus sign in front of the t we can see that as t increases x must decrease (we can verify with a quick derivative/Calculus I analysis if we want to). This means that the graph must be tracing out from right to left as the table of values above in the table also indicates.

Using a quick Calculus analysis of one, or both, of the parametric equations is often a better and easier method for determining the direction of motion for a parametric curve. For “simple” parametric equations we can often get the direction based on a quick glance at the parametric equations and it avoids having to pick “nice” values of t for a table.

Step 4

Let's now get the limits on x and y and note that we really do need these before we start sketching the curve!

In this case we have a parabola that opens downward and we could use that to get a general set of limits on x and y . However, for this problem we've also restricted the values of t that we're using and that will in turn restrict the values of x and y that we can use for the sketch of the graph.

As we discussed above we know that the graph will sketch out from right to left and so the rightmost value of x will come from $t = 0$, which is $x = 4$. Likewise, the leftmost value of y will come from $t = 3$, which is $x = -2$. So, from this we can see the limits on x must be $-2 \leq x \leq 4$.

For the limits on the y we've got to be a little more careful. First, we know that the vertex occurs in the given range of t 's and because the parabola opens downward the largest value of y we will have is

$y = \frac{21}{4}$, i.e. the y -coordinate of the vertex. Also, because the parabola opens downward we know that the smallest value of y will have to be at one of the endpoints. So, for $t = 0$ we have $y = 3$ and for $t = 3$ we have $y = -15$. Therefore, the limits on y must be $-15 \leq y \leq \frac{21}{4}$.


So, putting all this together here are the limits on x and y .

$$-2 < x < 4 \qquad -15 \leq y \leq \frac{21}{4}$$

Note that for this problem we must have these limits prior to the sketching step. Because we've restricted the values of t to use we will have limits on x and y (as we just discussed) and so we will only have a portion of the graph of the full parabola. Having these limits will allow us to get the sketch of the parametric curve.

Step 5

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included the points from our table because we had them but we won't always include them as we are often only interested in the sketch itself and the direction of motion.

Also note that it is vitally important that we not extend the graph past the $t = 0$ and $t = 3$ points. If we extend the graph past these points we are implying that the graph will extend past them and of course it doesn't!

3. Eliminate the parameter for the following set of parametric equations, sketch the graph of the parametric curve and give any limits that might exist on x and y .

$$x = \sqrt{t+1} \qquad y = \frac{1}{t+1} \qquad t > -1$$

Step 1

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations that is actually really easy to do if we notice the following.

$$x = \sqrt{t+1} \quad \Rightarrow \quad x^2 = t+1$$

With this we can quickly convert the y equation to,

$$y = \frac{1}{x^2}$$

Step 2

At this point we can get limits on x and y pretty quickly so let's do that.

First, we know that square roots always return positive values (or zero of course) and so from the x equation we see that we must have $x > 0$. Note as well that this must be a strict inequality because the inequality restricting the range of t 's is also a strict inequality. In other words, because we aren't allowing $t = -1$ we will never get $x = 0$.

Speaking of which, you do see why we've restricted the t 's don't you?

Now, from our restriction on t we know that $t + 1 > 0$ and so from the y parametric equation we can see that we also must have $y > 0$. This matches what we see from the equation without the parameter we found in Step 1.

So, putting all this together here are the limits on x and y .

$$x > 0 \quad y > 0$$

Note that for this problem these limits are important (or at least the x limits are important). Because of the x limit we get from the parametric equation we can see that we won't have the full graph of the equation we found in the first step. All we will have is the portion that corresponds to $x > 0$.

Step 3

Before we sketch the graph of the parametric curve recall that all parametric curves have a direction of motion, *i.e.* the direction indicating increasing values of the parameter, t in this case.

There are several ways to get the direction of motion for the curve. One is to plug in values of t into the parametric equations to get some points that we can use to identify the direction of motion.

Here is a table of values for this set of parametric equations.

t	x	y
-0.95	0.2236	20
-0.75	0.5	4
0	1	1
2	$\sqrt{3}$	$\frac{1}{3}$

Note that there is an easier way (probably – it will depend on you of course) to determine direction of motion. Take a quick look at the x equation.

$$x = \sqrt{t+1}$$

Increasing the value of t will also cause $t + 1$ to increase and the square root will also increase (we can verify with a quick derivative/Calculus I analysis if we want to). This means that the graph must be tracing out from left to right as the table of values above in the table supports.

Likewise, we could use the y equation.


$$y = \frac{1}{t+1}$$

Again, we know that as t increases so does $t + 1$. Because the $t + 1$ is in the denominator we can further see that increasing this will cause the fraction, and hence y , to decrease. This means that the graph must be tracing out from top to bottom as both the x equation and table of values supports.

Using a quick Calculus analysis of one, or both, of the parametric equations is often a better and easier method for determining the direction of motion for a parametric curve. For “simple” parametric equations we can often get the direction based on a quick glance at the parametric equations and it avoids having to pick “nice” values of t for a table.

Step 4

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included the points from our table because we had them but we won't always include them as we are often only interested in the sketch itself and the direction of motion.

4. Eliminate the parameter for the following set of parametric equations, sketch the graph of the parametric curve and give any limits that might exist on x and y .

$$x = 3\sin(t) \quad y = -4\cos(t) \quad 0 \leq t \leq 2\pi$$

Step 1

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we will make use of the well-known trig identity,

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

We can solve each of the parametric equations for sine and cosine as follows,

$$\sin(t) = \frac{x}{3} \quad \cos(t) = -\frac{y}{4}$$

Plugging these into the trig identity gives,

$$\left(-\frac{y}{4}\right)^2 + \left(\frac{x}{3}\right)^2 = 1 \quad \Rightarrow \quad \frac{x^2}{9} + \frac{y^2}{16} = 1$$

Therefore, the parametric curve will be some or all of the ellipse above.

We have to be careful when eliminating the parameter from a set of parametric equations. The graph of the resulting equation in only x and y may or may not be the graph of the parametric curve. Often, although not always, the parametric curve will only be a portion of the curve from the equation in terms of only x and y . Another situation that can happen is that the parametric curve will retrace some or all of the curve from the equation in terms of only x and y more than once.

The next few steps will help us to determine just how much of the ellipse we have and if it retraces the ellipse, or a portion of the ellipse, more than once.

Before we proceed with the rest of the problem let's first note that there is really no set order for doing the steps. They can often be done in different orders and in some cases may actually be easier to do in different orders. The order we'll be following here is used simply because it is the order that I'm used to working them in. If you find a different order would be best for you then that is the order you should use.

Step 2

At this point we can get a good idea on what the limits on x and y are going to be so let's do that. Note that often we won't get the actual limits on x and y in this step. All we are really finding here is the largest possible range of limits for x and y . Having these can sometimes be useful for later steps and so we'll get them here.

We can use our knowledge of sine and cosine to get the following inequalities. Note as discussed above however that these may not be the limits on x and y we are after.

$$-1 \leq \sin(t) \leq 1$$

$$-3 \leq 3\sin(t) \leq 3$$

$$-3 \leq x \leq 3$$

$$-1 \leq \cos(t) \leq 1$$

$$4 \geq -4\cos(t) \geq -4$$

$$-4 \leq y \leq 4$$

Note that to find these limits in general we just start with the appropriate trig function and then build up the equation for x and y by first multiplying the trig function by any coefficient, if present, and then adding/subtracting any numbers that might be present (not needed in this case). This, in turn, gives us the largest possible set of limits for x and y . Just remember to be careful when multiplying an inequality by a negative number. Don't forget to flip the direction of the inequalities when doing this.

Now, at this point we need to be a little careful. What we've actually found here are the largest possible inequalities for the limits on x and y . This set of inequalities for the limits on x and y assume that the parametric curve will be completely traced out at least once for the range of t 's we were given in the problem statement. It is always possible that the curve will not trace out a full trace in the given range of t 's. In a later step we'll determine if the parametric curve does trace out a full trace and hence determine the actual limits on x and y .

Before we move onto the next step there are a couple of issues we should quickly discuss.

First, remember that when we talk about the parametric curve tracing out once we are not necessarily talking about the ellipse itself being fully traced out. The parametric curve will be at most the full ellipse and we haven't determined just yet how much of the ellipse the parametric curve will trace out. So, one trace of the parametric curve refers to the largest portion of the ellipse that the parametric curve can possibly trace out given no restrictions on t .

Second, if we can't completely determine the actual limits on x and y at this point why did we do them here? In part we did them here because we can and the answer to this step often does end up being the limits on x and y . Also, there are times where knowing the largest possible limits on x and/or y will be convenient for some of the later steps.

Finally, we can sometimes get these limits from the sketch of the parametric curve. However, there are some parametric equations that we can't easily get the sketch without doing this step. We'll eventually do some problems like that.

Step 3

Before we sketch the graph of the parametric curve recall that all parametric curves have a direction of motion, *i.e.* the direction indicating increasing values of the parameter, t in this case.

There are several ways to get the direction of motion for the curve. One is to plug in values of t into the parametric equations to get some points that we can use to identify the direction of motion.

Here is a table of values for this set of parametric equations. In this case we were also given a range of t 's and we need to restrict the t 's in our table to that range.

t	x	y
0	0	-4
$\frac{\pi}{2}$	3	0
π	0	4
$\frac{3\pi}{2}$	-3	0
2π	0	-4

Now, this table seems to suggest that the parametric equation will follow the ellipse in a counter clockwise rotation. It also seems to suggest that the ellipse will be traced out exactly once.

However, tables of values for parametric equations involving sine and/or cosine equations can be deceptive.

Because sine and cosine oscillate it is possible to choose “bad” values of t that suggest a single trace when in fact the curve is tracing out faster than we realize and it is in fact tracing out more than once. We’ll need to do some extra analysis to verify if the ellipse traces out once or more than once.

Also, just because the table suggests a particular direction doesn’t actually mean it is going in that direction. It could be moving in the opposite direction at a speed that just happens to match the points you got in the table. Go back to the notes and check out Example 5. Plug in the points we used in our table above and you’ll get a set of points that suggest the curve is tracing out clockwise when in fact it is tracing out counter clockwise!

Note that because this is such a “bad” way of getting the direction of motion we put it in its own step so we could discuss it in detail. The actual method we’ll be using is in the next step and we’ll not be doing table work again unless it is absolutely required for some other part of the problem.

Step 4

As suggested in the previous step the table of values is not a good way to get direction of motion for parametric curves involving trig function so let’s go through a much better way of determining the direction of motion. This method takes a little time to think things through but it will always get the correct direction if you take the time.

First, let’s think about what happens if we start at $t = 0$ and increase t to $t = \pi$.

As we cover this range of t ’s we know that cosine starts at 1, decreasing through zero and finally stops at -1. So, that means that y will start at $y = -4$ (*i.e.* where cosine is 1), go through the x -axis (*i.e.* where cosine is zero) and finally stop at $y = 4$ (*i.e.* where cosine is -1). Now, this doesn’t give us a direction of motion as all it really tells us that y increases and it could do this following the right side of the ellipse (*i.e.* counter clockwise) or it could do this following the left side of the ellipse (*i.e.* clockwise).

So, let’s see what the behavior of sine in this range tells us. Starting at $t = 0$ we know that sine will be zero and so x will also be zero. As t increases to $t = \frac{\pi}{2}$ we know that sine increases from zero to one and so x will increase from zero to three. Finally, as we further increase t to $t = \pi$ sine will decrease from one back to zero and so x will also decrease from three to zero.

So, taking the x and y analysis above together we can see that at $t = 0$ the curve will start at the point $(0, -4)$. As we increase t to $t = \frac{\pi}{2}$ the curve will have to follow the ellipse with increasing x and y until it hits the point $(3, 0)$. The only way we can reach this second point and have the correct increasing behavior for both x and y is to move in a counter clockwise direction along the right half of the ellipse.

If we further increase t from $t = \frac{\pi}{2}$ to $t = \pi$ we can see that y must continue to increase but x now decreases until we get to the point $(0, 4)$ and again the only way we can reach this third point and have the required increasing/decreasing information for y/x respectively is to be moving in a counter clockwise direction along the right half.

We can do a similar analysis increasing t from $t = \pi$ to $t = 2\pi$ to see that we must still move in a counter clockwise direction that takes us through the point $(-3, 0)$ and then finally ending at the point $(0, -4)$.

So, from this analysis we can see that the curve must be tracing out in a counter clockwise direction.

This analysis seems complicated and maybe not so easy to do the first few times you see it. However, once you do it a couple of times you'll see that it's not quite as bad as it initially seems to be. Also, it really is the only way to guarantee that you've got the correct direction of motion for the curve when dealing with parametric equations involving sine and/or cosine.

If you had trouble visualizing how sine and cosine changed as we increased t you might want to do a quick sketch of the graphs of sine and cosine and you'll see right away that we were correct in our analysis of their behavior as we increased t .

Step 5

Okay, in the last step notice that we also showed that the curve will trace the ellipse out exactly once in the given range of t 's. However, let's assume that we hadn't done the direction analysis yet and see if we can determine this without the direction analysis.

This is actually pretty simple to do, or at least simpler than the direction analysis. All it requires is that you know where sine and cosine are zero, 1 and -1. If you recall your unit circle it's always easy to know where sine and cosine have these values. We'll also be able to verify the ranges of x and y found in Step 2 were in fact the actual ranges for x and y .

Let's start with the "initial" point on the curve, *i.e.* the point at the left end of our range of t 's, $t = 0$ in this case. Where you start this analysis is really dependent upon the set of parametric equations, the parametric curve and/or if there is a range of t 's given. Good starting points are the "initial" point, one of the end points of the curve itself (if the curve does have endpoints) or $t = 0$. Sometimes one option will be better than the others and other times it won't matter.

In this case two of the options are the same point so it seems like a good point to use.

So, at $t = 0$ we are at the point $(0, -4)$. We know that the parametric curve is some or all of the ellipse we found in the first step. So, at this point let's assume it is the full ellipse and ask ourselves the following question. When do we get back to this point? Or, in other words, what is the next value of t after $t = 0$ (since that is the point we choose to start off with) are we back at the point $(0, -4)$?

Before doing this let's quickly note that if the parametric curve doesn't get back to this point we'll determine that in the following analysis and that will be useful in helping us to determine how much of the ellipse will get traced out by the parametric curve.

Okay let's back to the analysis. In order to be at the point $(0, -4)$ we know we must have $\sin(t) = 0$ (only way to get $x = 0$!) and we must have $\cos(t) = 1$ (only way to get $y = -4$!). For $t > 0$ we know that $\sin(t) = 0$ at $t = \pi, 2\pi, 3\pi, \dots$ and likewise we know that $\cos(t) = 1$ at $t = 2\pi, 4\pi, 6\pi, \dots$. The first value of t that is in both lists is $t = 2\pi$ and so this is the next value of t that will put us at that point.

This tells us several things. First, we found that the parametric equation will get back to the initial point and so it is possible for the parametric equation to trace out the full ellipse.


Secondly, we got back to the point $(0, -4)$ at the very last t from the range of t 's we were given in the problem statement and so the parametric curve will trace out the ellipse exactly once for the given range of t 's.

Finally, from this analysis we found the parametric curve traced out the full ellipse in the range of t 's given in the problem statement and so we know now that the limits of x and y we found in Step 2 are in fact the actual limits on x and y for this curve.

As a final comment from this step let's note that this analysis in this step was a little easier than normal because the argument of the trig functions was just a t as opposed to say $2t$ or $\frac{1}{3}t$ which does make the analysis a tiny bit more complicated. We'll see how to deal with these kinds of arguments in the next couple of problems.

Step 6

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included the points from our table because we had them but we won't always include them as we are often only interested in the sketch itself and the direction of motion.

Also, because the problem asked for it here are the formal limits on x and y for this parametric curve.

$$-3 \leq x \leq 3 \qquad -4 \leq y \leq 4$$

As a final set of thoughts for this problem you really should go back and make sure you understand the processes we went through in Step 4 and Step 5. Those are often the best way of getting at the information we found in those steps. The processes can seem a little mysterious at first but once you've done a couple you'll find it isn't as bad as they might have first appeared.

Also, for the rest of the problems in this section we'll build a table of t values only if it is absolutely necessary for the problem. In other words, the process we used in Step 4 and 5 will be the processes we'll be using to get direction of motion for the parametric curve and to determine if the curve is traced out more than once or not.

You should also take a look at problems 5 and 6 in this section and contrast the number of traces of the curve with this problem. The only difference in the set of parametric equations in problems 4, 5 and 6 is the argument of the trig functions. After going through these three problems can you reach any conclusions on how the argument of the trig functions will affect the parametric curves for this type of parametric equations?

5. Eliminate the parameter for the following set of parametric equations, sketch the graph of the parametric curve and give any limits that might exist on x and y .

$$x = 3 \sin(2t) \quad y = -4 \cos(2t) \quad 0 \leq t \leq 2\pi$$

Step 1

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we will make use of the well-known trig identity,

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

We can solve each of the parametric equations for sine and cosine as follows,

$$\sin(2t) = \frac{x}{3} \quad \cos(2t) = -\frac{y}{4}$$

Plugging these into the trig identity (remember the identity holds as long as the argument of both trig functions, $2t$ in this case, is the same) gives,

$$\left(-\frac{y}{4}\right)^2 + \left(\frac{x}{3}\right)^2 = 1 \quad \Rightarrow \quad \frac{x^2}{9} + \frac{y^2}{16} = 1$$

Therefore, the parametric curve will be some or all of the ellipse above.

We have to be careful when eliminating the parameter from a set of parametric equations. The graph of the resulting equation in only x and y may or may not be the graph of the parametric curve. Often, although not always, the parametric curve will only be a portion of the curve from the equation in terms of only x and y . Another situation that can happen is that the parametric curve will retrace some or all of the curve from the equation in terms of only x and y more than once.

This observation is especially important for this problem. The next few steps will help us to determine just how much of the ellipse we have and if it retraces the ellipse, or a portion of the ellipse, more than once.

Before we proceed with the rest of the problem let's first note that there is really no set order for doing the steps. They can often be done in different orders and in some cases may actually be easier to do in different orders. The order we'll be following here is used simply because it is the order that I'm used to working them in. If you find a different order would be best for you then that is the order you should use.

Step 2

At this point we can get a good idea on what the limits on x and y are going to be so let's do that. Note that often we won't get the actual limits on x and y in this step. All we are really finding here is the largest possible range of limits for x and y . Having these can sometimes be useful for later steps and so we'll get them here.

We can use our knowledge of sine and cosine to determine the limits on x and y as follows,

$$\begin{array}{ll} -1 \leq \sin(2t) \leq 1 & -1 \leq \cos(2t) \leq 1 \\ -3 \leq 3\sin(2t) \leq 3 & 4 \geq -4\cos(2t) \geq -4 \\ -3 \leq x \leq 3 & -4 \leq y \leq 4 \end{array}$$

Note that to find these limits in general we just start with the appropriate trig function and then build up the equation for x and y by first multiplying the trig function by any coefficient, if present, and then adding/subtracting any numbers that might be present (not needed in this case). This, in turn, gives us the largest possible set of limits for x and y . Just remember to be careful when multiplying an inequality by a negative number. Don't forget to flip the direction of the inequalities when doing this.

Now, at this point we need to be a little careful. What we've actually found here are the largest possible inequalities for the limits on x and y . This set of inequalities for the limits on x and y assume that the parametric curve will be completely traced out at least once for the range of t 's we were given in the problem statement. It is always possible that the curve will not trace out a full trace in the given range of t 's. In a later step we'll determine if the parametric curve does trace out a full trace and hence determine the actual limits on x and y .

Before we move onto the next step there are a couple of issues we should quickly discuss.

First, remember that when we talk about the parametric curve tracing out once we are not necessarily talking about the ellipse itself being fully traced out. The parametric curve will be at most the full ellipse and we haven't determined just yet how much of the ellipse the parametric curve will trace out. So, one trace of the parametric curve refers to the largest portion of the ellipse that the parametric curve can possibly trace out given no restrictions on t .

Second, if we can't completely determine the actual limits on x and y at this point why did we do them here? In part we did them here because we can and the answer to this step often does end up being the limits on x and y . Also, there are times where knowing the largest possible limits on x and/or y will be convenient for some of the later steps.

Finally, we can sometimes get these limits from the sketch of the parametric curve. However, there are some parametric equations that we can't easily get the sketch without doing this step. We'll eventually do some problems like that.

Step 3

Before we sketch the graph of the parametric curve recall that all parametric curves have a direction of motion, *i.e.* the direction indicating increasing values of the parameter, t in this case.

In previous problems one method we looked at was to build a table of values for a sampling of t 's in the range provided. However, as we discussed in Problem 4 of this section tables of values for parametric equations involving trig functions they can be deceptive and so we aren't going to use them to determine the direction of motion for this problem.

Also, as noted in the discussion in Problem 4 it also might help to have the graph of sine and cosine handy to look at since we'll be talking a lot about the behavior of sine/cosine as we increase the argument.

So for this problem we'll just do the analysis of the behavior of sine and cosine in the range of t 's we were provided to determine the direction of motion. We'll be doing a quicker version of the analysis here than we did in Problem 4 so you might want to go back and check that problem out if you have trouble following everything we're going here.

Let's start at $t = 0$ since that is the first value of t in the range of t 's we were given in the problem. This means we'll be starting the parametric curve at the point $(0, -4)$.

Now, what happens if we start to increase t ? First, if we increase t then we also increase $2t$, the argument of the trig functions in the parametric equations. So, what does this mean for $\sin(2t)$ and $\cos(2t)$? Well initially, we know that $\sin(2t)$ will increase from zero to one and at the same time $\cos(2t)$ will also have to decrease from one to zero.

So, this means that x (given by $x = 3\sin(2t)$) will have to increase from 0 to 3. Likewise, it means that y (given by $y = -4\cos(2t)$) will have to increase from -4 to 0. For the y equation note that while the cosine is decreasing the minus sign on the coefficient means that y itself will actually be increasing.

Because this behavior for x and y must be happening at simultaneously we can see that the only possibility is for the parametric curve to start at $(0, -4)$ and as we increase the value of t we must move to the right in the counter clockwise direction until we reach the point $(3, 0)$.

Okay, we're now at the point $(3, 0)$, so $\sin(2t) = 1$ and $\cos(2t) = 0$. Let's continue to increase t . A further increase of t will force $\sin(2t)$ to decrease from 1 to 0 and at the same time $\cos(2t)$ will decrease from 0 to -1.

In terms of x and y this means that, at the same time, x will now decrease from 3 to 0 while y will continue to increase from 0 to 4 (again the minus sign on the y equation means y must increase as the cosine decreases from 0 to -1). So, we must be continuing to move in a counter clockwise direction until we reach the point $(0, 4)$.

For the remainder we'll go a little quicker in the analysis and just discuss the behavior of x and y and skip the discussion of the behavior of the sine and cosine.

Another increase in t will force x to decrease from 0 to -3 and at the same time y will have to also decrease from 4 to 0. The only way for this to happen simultaneously is to move along the ellipse starting that $(0, 4)$ in a counter clockwise motion until we reach $(-3, 0)$.

Continuing to increase t and we can see that, at the same time, x will increase from -3 to 0 and y will decrease from 0 to -4 . Or, in other words we're moving along the ellipse in a counter clockwise motion from $(-3, 0)$ to $(0, -4)$.

At this point we've gotten back to the starting point and we got back to that point by always going in a counter clockwise direction and did not retrace any portion of the graph and so we can now safely say that the direction of motion for this curve will always counter clockwise.

We have to be very careful here to continue the analysis until we get back to the starting point and see just how we got back there. It is possible, as we'll see in later problems, for us to get back there by retracing back over the curve. This will have an effect on the direction of motion for the curve (*i.e.* the direction will change!). In this case however since we got back to the starting point without retracing any portion of the curve we know the direction will remain counter clockwise.

Step 4

Let's now think about how much of the ellipse is actually traced out or if the ellipse is traced out more than once for the range of t 's we were given in the problem. We'll also be able to verify if the ranges of x and y we found in Step 2 are the correct ones or if we need to modify them (and we'll also determine just how to modify them if we need to).

Be careful to not draw any conclusions about how much of the ellipse is traced out from the analysis in the previous step. If we follow that analysis we see a full single trace of the ellipse. However, we didn't ever really mention any values of t with the exception of the starting value. Because of that we can't really use the analysis in the previous step to determine anything about how much of the ellipse we trace out or how many times we trace the ellipse out.

Let's go ahead and start this portion out at the same value of t we started with in the previous step. So, at $t = 0$ we are at the point $(0, -4)$. Now, when do we get back to this point? Or, in other words, what is the next value of t after $t = 0$ (since that is the point we choose to start off with) are we at the point $(0, -4)$?

In order to be at this point we know we must have $\sin(2t) = 0$ (only way to get $x = 0$!) and we must have $\cos(2t) = 1$ (only way to get $y = -4$!). Note the arguments of the sine and cosine! That is very important for this step.

Now, for $t > 0$ we know that $\sin(2t) = 0$ at $2t = \pi, 2\pi, 3\pi, \dots$ and likewise we know that $\cos(2t) = 1$ at $2t = 2\pi, 4\pi, 6\pi, \dots$. Again, note the arguments of sine and cosine here! Because we want $\sin(2t)$ and $\cos(2t)$ to have certain values we need to determine the values of $2t$ we need to achieve the values of sine and cosine that we are looking for.

The first value of $2t$ that is in both lists is $2t = 2\pi$. This now tells us the value of t we need to get back to the starting point. We just need to solve this for t !

$$2t = 2\pi \quad \Rightarrow \quad t = \pi$$

So, we will get back to the starting point, without retracing any portion of the ellipse, important in some later problems, when we reach $t = \pi$.

But this is in the middle of the range of t 's we were given! So, just what does this mean for us? Well first of all, provided the argument of the sine/cosine is only in terms of t , as opposed to t^2 or \sqrt{t} for example, the "net" range of t 's for one trace will always be the same. So, we got one trace in the range of $0 \leq t \leq \pi$ and so the "net" range of t 's here is $\pi - 0 = \pi$ and so any range of t 's that span π will trace out the ellipse exactly once.


This means that the ellipse will also trace out exactly once in the range $\pi \leq t \leq 2\pi$. So, in this case, it looks like the ellipse will be traced out twice in the range $0 \leq t \leq 2\pi$.

This analysis also has shown us that the parametric curve traces out the full ellipse in the range of t 's given in the problem statement (more than once in fact!) and so we know now that the limits of x and y we found in Step 2 are in fact the actual limits on x and y for this curve.

Before we leave this step we should note that once you get pretty good at the direction analysis we did in Step 3 you can combine the analysis Steps 3 and 4 into a single step to get both the direction and portion of the curve that is traced out. Initially however you might find them a little easier to do them separately.

Step 5

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included a set of t 's to illustrate a handful of points and their corresponding values of t 's. For some practice you might want to follow the analysis from Step 4 to see if you can verify the values of t for the other three points on the graph. It would, of course, be easier to just plug them in to verify, but the practice of the process of the Step 4 analysis might be useful to you.

Also, because the problem asked for it here are the formal limits on x and y for this parametric curve.

$$-3 \leq x \leq 3 \qquad -4 \leq y \leq 4$$

You should also take a look at problems 4 and 6 in this section and contrast the number of traces of the curve with this problem. The only difference in the set of parametric equations in problems 4, 5 and 6 is the argument of the trig functions. After going through these three problems can you reach any conclusions on how the argument of the trig functions will affect the parametric curves for this type of parametric equations?

6. Eliminate the parameter for the following set of parametric equations, sketch the graph of the parametric curve and give any limits that might exist on x and y .

$$x = 3 \sin\left(\frac{1}{3}t\right) \quad y = -4 \cos\left(\frac{1}{3}t\right) \quad 0 \leq t \leq 2\pi$$

Step 1

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we will make use of the well-known trig identity,

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

We can solve each of the parametric equations for sine and cosine as follows,

$$\sin\left(\frac{1}{3}t\right) = \frac{x}{3} \qquad \cos\left(\frac{1}{3}t\right) = -\frac{y}{4}$$

Plugging these into the trig identity (remember the identity holds as long as the argument of both trig functions, $\frac{1}{3}t$ in this case, is the same) gives,

$$\left(-\frac{y}{4}\right)^2 + \left(\frac{x}{3}\right)^2 = 1 \qquad \Rightarrow \qquad \frac{x^2}{9} + \frac{y^2}{16} = 1$$

Therefore, the parametric curve will be some or all of the ellipse above.

We have to be careful when eliminating the parameter from a set of parametric equations. The graph of the resulting equation in only x and y may or may not be the graph of the parametric curve. Often, although not always, the parametric curve will only be a portion of the curve from the equation in terms

of only x and y . Another situation that can happen is that the parametric curve will retrace some or all of the curve from the equation in terms of only x and y more than once.

This observation is especially important for this problem. The next few steps will help us to determine just how much of the ellipse we have and if it retraces the ellipse, or a portion of the ellipse, more than once.

Before we proceed with the rest of the problem let's first note that there is really no set order for doing the steps. They can often be done in different orders and in some cases may actually be easier to do in different orders. The order we'll be following here is used simply because it is the order that I'm used to working them in. If you find a different order would be best for you then that is the order you should use.

Step 2

At this point we can get a good idea on what the limits on x and y are going to be so let's do that. Note that often we won't get the actual limits on x and y in this step. All we are really finding here is the largest possible range of limits for x and y . Having these can sometimes be useful for later steps and so we'll get them here.

We can use our knowledge of sine and cosine to determine the limits on x and y as follows,

$$\begin{array}{ll} -1 \leq \sin\left(\frac{1}{3}t\right) \leq 1 & -1 \leq \cos\left(\frac{1}{3}t\right) \leq 1 \\ -3 \leq 3\sin\left(\frac{1}{3}t\right) \leq 3 & 4 \geq -4\cos\left(\frac{1}{3}t\right) \geq -4 \\ -3 \leq x \leq 3 & -4 \leq y \leq 4 \end{array}$$

Note that to find these limits in general we just start with the appropriate trig function and then build up the equation for x and y by first multiplying the trig function by any coefficient, if present, and then adding/subtracting any numbers that might be present (not needed in this case). This, in turn, gives us the largest possible set of limits for x and y . Just remember to be careful when multiplying an inequality by a negative number. Don't forget to flip the direction of the inequalities when doing this.

Now, at this point we need to be a little careful. What we've actually found here are the largest possible inequalities for the limits on x and y . This set of inequalities for the limits on x and y assume that the parametric curve will be completely traced out at least once for the range of t 's we were given in the problem statement. It is always possible that the curve will not trace out a full trace in the given range of t 's. In a later step we'll determine if the parametric curve does trace out a full trace and hence determine the actual limits on x and y .

Before we move onto the next step there are a couple of issues we should quickly discuss.

First, remember that when we talk about the parametric curve tracing out once we are not necessarily talking about the ellipse itself being fully traced out. The parametric curve will be at most the full ellipse and we haven't determined just yet how much of the ellipse the parametric curve will trace out. So, one trace of the parametric curve refers to the largest portion of the ellipse that the parametric curve can possibly trace out given no restrictions on t . This is especially important for this problem!

Second, if we can't completely determine the actual limits on x and y at this point why did we do them here? In part we did them here because we can and the answer to this step often does end up being the limits on x and y . Also, there are times where knowing the largest possible limits on x and/or y will be convenient for some of the later steps.

Finally, we can sometimes get these limits from the sketch of the parametric curve. However, there are some parametric equations that we can't easily get the sketch without doing this step. We'll eventually do some problems like that.

Step 3

Before we sketch the graph of the parametric curve recall that all parametric curves have a direction of motion, *i.e.* the direction indicating increasing values of the parameter, t in this case.

In previous problems one method we looked at was to build a table of values for a sampling of t 's in the range provided. However, as we discussed in Problem 4 of this section tables of values for parametric equations involving trig functions they can be deceptive and so we aren't going to use them to determine the direction of motion for this problem.

Also, as noted in the discussion in Problem 4 it also might help to have the graph of sine and cosine handy to look at since we'll be talking a lot about the behavior of sine/cosine as we increase the argument.

So for this problem we'll just do the analysis of the behavior of sine and cosine in the range of t 's we were provided to determine the direction of motion. We'll be doing a quicker version of the analysis here than we did in Problem 4 so you might want to go back and check that problem out if you have trouble following everything we're going here.

Let's start at $t = 0$ since that is the first value of t in the range of t 's we were given in the problem. This means we'll be starting the parametric curve at the point $(0, -4)$.

Now, what happens if we start to increase t ? First, if we increase t then we also increase $\frac{1}{3}t$, the argument of the trig functions in the parametric equations. So, what does this mean for $\sin\left(\frac{1}{3}t\right)$ and $\cos\left(\frac{1}{3}t\right)$? Well initially, we know that $\sin\left(\frac{1}{3}t\right)$ will increase from zero to one and at the same time $\cos\left(\frac{1}{3}t\right)$ will also have to decrease from one to zero.

So, this means that x (given by $x = 3\sin\left(\frac{1}{3}t\right)$) will have to increase from 0 to 3. Likewise, it means that y (given by $y = -4\cos\left(\frac{1}{3}t\right)$) will have to increase from -4 to 0. For the y equation note that while the cosine is decreasing the minus sign on the coefficient means that y itself will actually be increasing.

Because this behavior for the x and y must be happening at simultaneously we can see that the only possibility is for the parametric curve to start at $(0, -4)$ and as we increase the value of t we must move to the right in the counter clockwise direction until we reach the point $(3, 0)$.

Okay, we're now at the point $(3, 0)$, so $\sin\left(\frac{1}{3}t\right) = 1$ and $\cos\left(\frac{1}{3}t\right) = 0$. Let's continue to increase t . A further increase of t will force $\sin\left(\frac{1}{3}t\right)$ to decrease from 1 to 0 and at the same time $\cos\left(\frac{1}{3}t\right)$ will decrease from 0 to -1.

In terms of x and y this means that, at the same time, x will now decrease from 3 to 0 while y will continue to increase from 0 to 4 (again the minus sign on the y equation means y must increase as the cosine decreases from 0 to -1). So, we must be continuing to move in a counter clockwise direction until we reach the point $(0, 4)$.

For the remainder we'll go a little quicker in the analysis and just discuss the behavior of x and y and skip the discussion of the behavior of the sine and cosine.

Another increase in t will force x to decrease from 0 to -3 and at the same time y will have to also decrease from 4 to 0. The only way for this to happen simultaneously is to move along the ellipse starting at $(0, 4)$ in a counter clockwise motion until we reach $(-3, 0)$.

Continuing to increase t and we can see that, at the same time, x will increase from -3 to 0 and y will decrease from 0 to -4. Or, in other words we're moving along the ellipse in a counter clockwise motion from $(-3, 0)$ to $(0, -4)$.

At this point we've gotten back to the starting point and we got back to that point by always going in a counter clockwise direction and did not retrace any portion of the graph and so we can now safely say that the direction of motion for this curve will always counter clockwise.

We have to be very careful here to continue the analysis until we get back to the starting point and see just how we got back there. It is possible, as we'll see in later problems, for us to get back there by retracing back over the curve. This will have an effect on the direction of motion for the curve (*i.e.* the direction will change!). In this case however since we got back to the starting point without retracing any portion of the curve we know the direction will remain counter clockwise.

Step 4

Let's now think about how much of the ellipse is actually traced out or if the ellipse is traced out more than once for the range of t 's we were given in the problem. We'll also be able to verify if the ranges of x and y we found in Step 2 are the correct ones or if we need to modify them (and we'll also determine just how to modify them if we need to).

Be careful to not draw any conclusions about how much of the ellipse is traced out from the analysis in the previous step. If we follow that analysis we see a full single trace of the ellipse. However, we didn't ever really mention any values of t with the exception of the starting value. Because of that we can't really use the analysis in the previous step to determine anything about how much of the ellipse we trace out or how many times we trace the ellipse out.

Let's go ahead and start this portion out at the same value of t we started with in the previous step. So, at $t = 0$ we are at the point $(0, -4)$. Now, when do we get back to this point? Or, in other words, what

is the next value of t after $t = 0$ (since that is the point we choose to start off with) are we at the point $(0, -4)$?

In order to be at this point we know we must have $\sin\left(\frac{1}{3}t\right) = 0$ (only way to get $x = 0$!) and we must have $\cos\left(\frac{1}{3}t\right) = 1$ (only way to get $y = -4$!). Note the arguments of the sine and cosine! That is very important for this step.

Now, for $t > 0$ we know that $\sin\left(\frac{1}{3}t\right) = 0$ at $\frac{1}{3}t = \pi, 2\pi, 3\pi, \dots$ and likewise we know that $\cos\left(\frac{1}{3}t\right) = 1$ at $\frac{1}{3}t = 2\pi, 4\pi, 6\pi, \dots$. Again, note the arguments of sine and cosine here! Because we want $\sin\left(\frac{1}{3}t\right)$ and $\cos\left(\frac{1}{3}t\right)$ to have certain values we need to determine the values of $\frac{1}{3}t$ we need to achieve the values of sine and cosine that we are looking for.

The first value of $\frac{1}{3}t$ that is in both lists is $\frac{1}{3}t = 2\pi$. This now tells us the value of t we need to get back to the starting point. We just need to solve this for t !

$$\frac{1}{3}t = 2\pi \quad \Rightarrow \quad t = 6\pi$$

So, we will get back to the starting point, without retracing any portion of the ellipse, important in some later problems, when we reach $t = 6\pi$.

At this point we have a problem that we didn't have in the previous two problems. We get back to the point $(0, -4)$ at $t = 6\pi$ and this is outside the range of t 's given in the problem statement, $0 \leq t \leq 2\pi$!

What this means for us is that the parametric curve will not trace out a full trace for the range of t 's we were given for this problem. It also means that the range of limits for x and y from Step 2 are not the correct limits for x and y .

We know from the Step 3 analysis that the parametric curve will trace out in a counter clockwise direction and from the analysis in this step it won't trace out a full trace.

So, we know the parametric curve will start when $t = 0$ at $(0, -4)$ and will trace out in a counter clockwise direction until $t = 2\pi$ at which we will be at the point,

$$\left(3 \sin\left(\frac{2\pi}{3}\right), -4 \cos\left(\frac{2\pi}{3}\right)\right) = \left(\frac{3\sqrt{3}}{2}, 2\right)$$


This "ending" point is in the first quadrant and so we know that the curve has to have passed through $(3, 0)$. This means that the limits on x are $0 \leq x \leq 3$. The limits on the y are simply those we get from the points $-4 \leq y \leq 2$.

Before we leave this step we should note that once you get pretty good at the direction analysis we did in Step 3 you can combine the analysis Steps 3 and 4 into a single step to get both the direction and

portion of the curve that is traced out. Initially however you might find them a little easier to do them separately.

Step 5

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included a set of t 's to illustrate a handful of points and their corresponding values of t 's. For some practice you might want to follow the analysis from Step 4 to see if you can verify the values of t for the other three points on the graph. It would, of course, be easier to just plug them in to verify, but the practice would of the Step 4 analysis might be useful to you.

Note as well that we included the full sketch of the ellipse as a dashed graph to help illustrate the portion of the ellipse that the parametric curve is actually covering.

Also, because the problem asked for it here are the formal limits on x and y for this parametric curve.

$$0 \leq x \leq 3 \qquad -4 \leq y \leq 2$$

You should also take a look at problems 4 and 5 in this section and contrast the number of traces of the curve with this problem. The only difference in the set of parametric equations in problems 4, 5 and 6 is the argument of the trig functions. After going through these three problems can you reach any conclusions on how the argument of the trig functions will affect the parametric curves for this type of parametric equations?

7. The path of a particle is given by the following set of parametric equations. Completely describe the path of the particle. To completely describe the path of the particle you will need to provide the following information.

(i) A sketch of the parametric curve (including direction of motion) based on the equation you get by eliminating the parameter.

(ii) Limits on x and y .

(iii) A range of t 's for a single trace of the parametric curve.

(iv) The number of traces of the curve the particle makes if an overall range of t 's is provided in the problem.

$$x = 3 - 2 \cos(3t) \quad y = 1 + 4 \sin(3t)$$

Step 1

There's a lot of information we'll need to find to fully answer this problem. However, for most of it we can follow the same basic ordering of steps we used for the first few problems in this section. We will need however to do a little extra work along the way.

Also, because most of the work here is similar to the work we did in Problems 4 – 6 of this section we won't be putting in as much explanation to a lot of the work we're doing here. So, if you need some explanation for some of the work you should go back to those problems and check the corresponding steps.

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we will make use of the well-known trig identity,

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

We can solve each of the parametric equations for sine and cosine as follows,

$$\cos(3t) = \frac{x-3}{-2} \quad \sin(3t) = \frac{y-1}{4}$$

Plugging these into the trig identity gives,

$$\left(\frac{x-3}{-2}\right)^2 + \left(\frac{y-1}{4}\right)^2 = 1 \quad \Rightarrow \quad \frac{(x-3)^2}{4} + \frac{(y-1)^2}{16} = 1$$

Therefore, the parametric curve will be some or all of the graph of this ellipse.

Step 2

At this point let's get our first guess as to the limits on x and y . As noted in previous problems what we're really finding here is the largest possible ranges for x and y . In later steps we'll determine if this the actual set of limits on x and y or if we have smaller ranges.

We can use our knowledge of sine and cosine to determine the limits on x and y as follows,

$$\begin{array}{ll}
 -1 \leq \cos(3t) \leq 1 & -1 \leq \sin(3t) \leq 1 \\
 2 \geq -2\cos(3t) \geq -2 & -4 \leq 4\sin(3t) \leq 4 \\
 5 \geq 3 - 2\cos(3t) \geq 1 & -3 \leq 1 + 4\sin(3t) \leq 5 \\
 1 \leq x \leq 5 & -3 \leq y \leq 5
 \end{array}$$

Remember that all we need to do is start with the appropriate trig function and then build up the equation for x and y by first multiplying the trig function by any coefficient, if present, and then adding/subtracting any numbers that might be present. We now have the largest possible set of limits for x and y .

This problem does not have a range of t 's that might restrict how much of the parametric curve gets sketched out. This means that the parametric curve will be fully traced out.

Remember that when we talk about the parametric curve getting fully traced out this doesn't, in general, mean the full ellipse we found in Step 1 gets traced out by the parametric equation. All "fully traced out" means, in general, is that whatever portion of the ellipse that is described by the set of parametric curves will be completely traced out.

However, for this problem let's also note as well that the ranges for x and y we found above also correspond the maximum ranges for x and y we get from the equation of the ellipse we found in Step 1. This means that, for this problem, the ellipse will get fully traced out at least once by the parametric curve and so these are the full limits on x and y .

Step 3

Let's next get the direction of motion for the parametric curve.

Let's use $t = 0$ as a "starting" point for this analysis. At $t = 0$ we are at the point $(1,1)$. If we increase t we can see that both x and y must increase until we get to the point $(3,5)$. Increasing t further from this point will force x to continue to increase, but y will now start to decrease until we reach the point $(5,1)$. Next, when we increase t further both x and y will decrease until we reach the point $(3,-3)$. Finally, increasing t even more we get x continuing to decrease while y starts to increase until we get back to $(1,1)$, the point we "started" the analysis at.

We didn't put a lot of "explanation into this but if you think about the parametric equations and how sine/cosine behave as you increase t you should see what's going on. In the x equation we see that the coefficient of the cosine is negative and so if cosine increases x must decrease and if cosine decreases x must increase. For the y equation the coefficient of the sine is positive and so both y and sine will increase or decrease at the same time.

Okay, in all of the analysis above we must be moving in a clockwise direction. Also, note that because of the oscillating nature of sine and cosine once we reach back to the "starting" point the behavior will simply repeat itself. This in turn tells us that once we arrive back at the "starting" point we will continue to trace out the parametric curve in a clockwise direction.

Step 4

From the analysis in the last step we saw that without any range of t 's restricting the parametric curve, which we don't have here, the parametric curve will completely trace out the ellipse that we found in Step 1.

Therefore, the next thing we should do is determine a range of t 's that it will take to complete one trace of the parametric curve. Note that one trace of the parametric curve means that no portion of the parametric curve will ever be retraced. For this problem that means we trace out the ellipse exactly once.

So, as with the last step let's "start" at the point $(1,1)$, which corresponds to $t = 0$. So, the next question to ask is what value of $t > 0$ will we reach this point again.

In order to be at the point $(1,1)$ we need to require that $\cos(3t) = 1$ and $\sin(3t) = 0$. So, for $t > 0$ we know we'll have $\cos(3t) = 1$ if $3t = 2\pi, 4\pi, 6\pi, \dots$ and we'll have $\sin(3t) = 0$ if $3t = \pi, 2\pi, 3\pi, \dots$.

The first value of t that is in both of these lists is $3t = 2\pi$. So, we'll get back to the "starting" point at,

$$3t = 2\pi \quad \Rightarrow \quad t = \frac{2\pi}{3}$$

Therefore, one trace will be completed in the range,

$$0 \leq t \leq \frac{2\pi}{3}$$

Note that this is only one possible answer here. Any range of t 's with a "net" range of $\frac{2\pi}{3}$ t 's, with the endpoints of the t range corresponding to start/end points of the parametric equation, will work. So, for example, any of the following ranges of t 's would also work.

$$-\frac{2\pi}{3} \leq t \leq 0 \qquad \frac{2\pi}{3} \leq t \leq \frac{4\pi}{3} \qquad -\frac{\pi}{3} \leq t \leq \frac{\pi}{3}$$


There are of course many other possible ranges of t 's for a one trace. Note however, as the last example above shows, because the full ellipse is traced out, each range doesn't all need to start/end at the same place. The range we originally arrived at as well as the first two ranges above all start/end at $(1,1)$ while the third range above starts/ends at $(5,1)$.

Step 5

Now that we have a range of t 's for one full trace of the parametric curve we could determine the number of traces the particle makes. However, because we weren't given an overall range of t 's we can't do that for this problem.

Step 6

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we included a set of t 's to illustrate where the particle is at while tracing out of the curve. For some practice you might want to follow the analysis from Step 4 to see if you can verify the values of t for the other three points on the graph. It would, of course, be easier to just plug them in to verify, but the practice would of the Step 4 analysis might be useful to you.

Here is also the formal answers for all the rest of the information that problem asked for.

Range of x :	$1 \leq x \leq 5$
Range of y :	$-3 \leq y \leq 5$
Range of t for one trace :	$0 \leq t \leq \frac{2\pi}{3}$
Total number of traces :	n/a

8. The path of a particle is given by the following set of parametric equations. Completely describe the path of the particle. To completely describe the path of the particle you will need to provide the following information.

(i) A sketch of the parametric curve (including direction of motion) based on the equation you get by eliminating the parameter.

(ii) Limits on x and y .

(iii) A range of t 's for a single trace of the parametric curve.

(iv) The number of traces of the curve the particle makes if an overall range of t 's is provided in the problem.

$$x = 4 \sin\left(\frac{1}{4}t\right) \quad y = 1 - 2 \cos^2\left(\frac{1}{4}t\right) \quad -52\pi \leq t \leq 34\pi$$

Step 1

There's a lot of information we'll need to find to fully answer this problem. However, for most of it we can follow the same basic ordering of steps we used for the first few problems in this section. We will need however to do a little extra work along the way.

Also, because most of the work here is similar to the work we did in Problems 4 – 6 of this section we won't be putting in as much explanation to a lot of the work we're doing here. So, if you need some explanation for some of the work you should go back to those problems and check the corresponding steps.

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations we will make use of the well-known trig identity,

$$\cos^2(\theta) + \sin^2(\theta) = 1$$

We can solve each of the parametric equations for sine and cosine as follows,

$$\sin\left(\frac{1}{4}t\right) = \frac{x}{4} \qquad \cos^2\left(\frac{1}{4}t\right) = \frac{y-1}{-2}$$

Plugging these into the trig identity gives,

$$\frac{y-1}{-2} + \left(\frac{x}{4}\right)^2 = 1 \qquad \Rightarrow \qquad y = \frac{x^2}{8} - 1$$

Therefore, with a little algebraic manipulation, we see that the parametric curve will be some or all of the parabola above. Note that while many parametric equations involving sines and cosines are some or all of an ellipse they won't all be as this problem shows. Do not get so locked into ellipses when seeing sines/cosines that you always just assume the curve will be an ellipse.

Step 2

At this point let's get our first guess as to the limits on x and y . As noted in previous problems what we're really finding here is the largest possible ranges for x and y . In later steps we'll determine if this the actual set of limits on x and y or if we have smaller ranges.

We can use our knowledge of sine and cosine to determine the limits on x and y as follows,

$$\begin{array}{ll} -1 \leq \sin\left(\frac{1}{4}t\right) \leq 1 & -1 \leq \cos\left(\frac{1}{4}t\right) \leq 1 \\ -4 \leq 4\sin\left(\frac{1}{4}t\right) \leq 4 & 0 \leq \cos^2\left(\frac{1}{4}t\right) \leq 1 \\ -4 \leq x \leq 4 & 0 \geq -2\cos^2\left(\frac{1}{4}t\right) \geq -2 \\ & 1 \geq 1 - 2\cos^2\left(\frac{1}{4}t\right) \geq -1 \\ & -1 \leq y \leq 1 \end{array}$$

Remember that all we need to do is start with the appropriate trig function and then build up the equation for x and y by first multiplying the trig function by any coefficient, if present, and then

adding/subtracting any numbers that might be present. We now have the largest possible set of limits for x and y .


Now, at this point we need to be a little careful. As noted above what we've actually found here are the largest possible ranges for the limits on x and y . This set of inequalities for the limits on x and y assume that the parametric curve will be fully traced out at least once for the range of t 's we were given in the problem statement. It is always possible that the parametric curve will not trace out a full trace in the given range of t 's. In a later step we'll determine if the parametric curve does trace out a full trace and hence determine the actual limits on x and y .

Remember that when we talk about the parametric curve getting fully traced out this doesn't, in general, mean the full parabola we found in Step 1 gets traced out by the parametric equation. All "fully traced out" means, in general, is that whatever portion of the parabola that is described by the set of parametric curves will be completely traced out.

In fact, for this problem, we can see that the parabola from Step 1 will not get fully traced out by the particle regardless of any range of t 's. The largest possible portion of the parabola that can be traced out by the particle is the portion that lies in the range of x and y given above. In a later step we'll determine if the largest possible portion of the parabola does get traced out or if the particle only traces out part of it.

Step 3

Let's next get the direction of motion for the parametric curve. For this analysis it might be useful to have a quick sketch of the largest possible parametric curve. So, here is a quick sketch of that.



The dashed line is the graph of the full parabola from Step 1 and the solid line is the portion that falls into our largest possible range of x and y we found in Step 2. As an aside here note that the two ranges are complimentary. In other words, if we sketch the graph only for the range of x we automatically get the range for y . Likewise, if we sketch the graph only for the range of y we automatically get the range for x . This is a good check for your graph. The x and y ranges should always match up!

When our parametric curve was an ellipse (the previous problem for example) no matter what point we started the analysis at the curve would eventually trace out around the ellipse and end up back at the starting point without ever going back over any portion of itself. The main issue we faced with the ellipse problem was we could rotate around the ellipse in a clockwise or a counter clockwise motion to

do this and a careful analysis of the behavior of both the x and y parametric equations was required to determine just which direction we were going.

With a parabola for our parametric curve things work a lot differently. Let's suppose that we "started" at the right end point (this is just randomly picked for no other reason that I'm right handed so don't think there is anything special about this point!) and it doesn't matter what t we use to get to that point.

At this point we know that we are at $x = 4$ and in order for x to have that value we must also have $\sin\left(\frac{1}{4}t\right) = 1$. Now, as we increase t from this point (again it doesn't matter just what the value of t is) the only option for sine is for it to decrease until it has the value $\sin\left(\frac{1}{4}t\right) = -1$. This in turn means that if we start at the right end point we have no option but to proceed along the curve going to the left.

However, we don't just reach the left end point and then stop! Once we are at $\sin\left(\frac{1}{4}t\right) = -1$ if we further increase t we know that sine will also increase until it has the value $\sin\left(\frac{1}{4}t\right) = 1$ and so we must move back along the curve to the right until we are back at the right end point.

Unlike the ellipse however, the only way for this to happen is for the particle to go back over the parabola moving in a rightward direction. Remember that the particle moves to the right or left it must trace out a portion of the parabola that we found in Step 1! Any particle traveling along the path given by the set of parametric equations must follow the graph of the parabola and never leave it.

In other words, if we don't put any restrictions on t a particle on this parametric curve will simply oscillate left and right along the portion of the parabola sketched out above. In this case however we do have a range of t 's so we'll need to determine a range of t 's for one trace to fully know the direction of motion information of the particle on this path and we'll do that in the next step. With a restriction on the range of t 's it is possible that the particle won't make a full trace or it might retrace some or all of the curve so we can't say anything definite about the direction of motion for the particle over the full range of t 's until the next step when we determine a range of t 's for one full trace of the curve.

Before we move on to the next step there is a quick topic we should address. We only used the x equation to do this analysis and never addressed the y -equation anywhere in the analysis. It doesn't really matter which one we use as both will give the same information.

Step 4

Now we need to determine a range of t 's for one full trace of the parametric curve. It is important for this step to remember that one full trace of the parametric curve means that no portion of the parametric curve can be retraced.

Note that one full trace does not mean that we get back to the "starting" point. When we dealt with an ellipse in the previous problem that was one trace because we did not need to retrace any portion of the ellipse to get back to the starting point. However, as we saw in the previous step that for our parabola here we would have to retrace the full curve to get back to the starting point.

So, one full trace of the parametric curve means we move from the right end point to the left end point only or visa-versa and move from the left end point to the right end point. Which direction we move

doesn't really matter here so let's get a range of t 's that take us from the left end point to the right end point.

In all the previous problems we've used $t = 0$ as our "starting" point but that won't work for this problem because that actually corresponds to the vertex of the parabola. We want to start at the left end point so the first part of this process is actually determine a t that will put us at the left end point.

In order to be at the left end point, $(-4,1)$, we need to require that $\sin\left(\frac{1}{4}t\right) = -1$ which occurs if $\frac{1}{4}t = \dots, -\frac{5\pi}{2}, -\frac{\pi}{2}, \frac{3\pi}{2}, \frac{7\pi}{2}, \dots$. We also need to require that $\cos\left(\frac{1}{4}t\right) = 0$ which occurs if $\frac{1}{4}t = \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$. There are going to be many numbers that are in both lists here so all we need to do is pick one and proceed. From the numbers that we've listed here we could use either $\frac{1}{4}t = -\frac{\pi}{2}$ or $\frac{1}{4}t = \frac{3\pi}{2}$. We'll use $\frac{1}{4}t = -\frac{\pi}{2}$, *i.e.* when $t = -2\pi$, simply because it is the first one that occurs in both lists. Therefore, we will be at the left end point when $t = -2\pi$.

Let's now move to the right end point, $(4,1)$. In order to get the range of t 's for one trace this means we'll need the next t with $t > -2\pi$ (which corresponds to $\frac{1}{4}t > -\frac{\pi}{2}$). To do this we need to require that $\sin\left(\frac{1}{4}t\right) = 1$ which occurs if $\frac{1}{4}t = \dots, -\frac{3\pi}{2}, \frac{\pi}{2}, \frac{5\pi}{2}, \frac{9\pi}{2}, \dots$ and we need to that $\cos\left(\frac{1}{4}t\right) = 0$ which occurs if $\frac{1}{4}t = \dots, -\frac{3\pi}{2}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{2}, \dots$.

The first t that is in both of these lists with $\frac{1}{4}t > -\frac{\pi}{2}$ is then $\frac{1}{4}t = \frac{\pi}{2}$, *i.e.* when $t = 2\pi$. So, the first t after $t = -2\pi$ that puts us at the right end point is $t = 2\pi$. This means that a range of t 's for one full trace of the parametric curve is then,

$$-2\pi \leq t \leq 2\pi$$

Note that this is only one possible answer here. Any range of t 's with a "net" range of $2\pi - (-2\pi) = 4\pi$ t 's, with the endpoints of the t range corresponding to start/end points of the parametric curve, will work. So, for example, any of the following ranges of t 's would also work.

$$-6\pi \leq t \leq -2\pi \qquad 2\pi \leq t \leq 6\pi \qquad 6\pi \leq t \leq 10\pi$$

The direction of motion for each may be different range of t 's of course. Some will trace out the curve moving from left to right while others will trace out the curve moving from right to left. Because the problem did not specify a particular direction any would work.

Note as well that the range $-2\pi \leq t \leq 2\pi$ falls completely inside the given range of t 's specified in the problem and so we know that the particle will trace out the curve more than once over the full range of t 's. Determining just how many times it traces over the curve will be determined in the next step.

Step 5


Now that we have a range of t 's for one full trace of the parametric curve we can determine the number of traces the particle makes.

This is a really easy step. We know the total time the particle was traveling and we know how long it takes for a single trace. Therefore,

$$\text{Number Traces} = \frac{\text{Total Time Traveled}}{\text{Time for One Trace}} = \frac{34\pi - (-52\pi)}{2\pi - (-2\pi)} = \frac{86\pi}{4\pi} = \frac{43}{2} = 21.5 \text{ traces}$$

Step 6

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we indicated the direction of motion by putting arrow heads going both directions in places on the curve. We also included a set of t 's for a couple of points to illustrate where the particle is at while tracing out of the curve. The dashed line is the continuation of the parabola from Step 1 to illustrate that our parametric curve is only a part of the parabola.

Here is also the formal answers for all the rest of the information that problem asked for.

Range of x :	$-4 \leq x \leq 4$
Range of y :	$-1 \leq y \leq 1$
Range of t for one trace :	$-2\pi \leq t \leq 2\pi$
Total number of traces :	21.5

9. The path of a particle is given by the following set of parametric equations. Completely describe the path of the particle. To completely describe the path of the particle you will need to provide the following information.

(i) A sketch of the parametric curve (including direction of motion) based on the equation you get by eliminating the parameter.

(ii) Limits on x and y .

(iii) A range of t 's for a single trace of the parametric curve.

(iv) The number of traces of the curve the particle makes if an overall range of t 's is provided in the problem.

$$x = \sqrt{4 + \cos\left(\frac{5}{2}t\right)} \quad y = 1 + \frac{1}{3}\cos\left(\frac{5}{2}t\right) \quad -48\pi \leq t \leq 2\pi$$

Step 1

There's a lot of information we'll need to find to fully answer this problem. However, for most of it we can follow the same basic ordering of steps we used for the first few problems in this section. We will need however to do a little extra work along the way.

Also, because most of the work here is similar to the work we did in Problems 4 – 6 of this section we won't be putting in as much explanation to a lot of the work we're doing here. So, if you need some explanation for some of the work you should go back to those problems and check the corresponding steps.

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations notice that we can quickly and easily eliminate the parameter simply by solving the y equation for cosine as follows,

$$\cos\left(\frac{5}{2}t\right) = 3y - 3$$

Plugging this into the cosine in the x equation gives,

$$x = \sqrt{4 + (3y - 3)} \quad \Rightarrow \quad x = \sqrt{1 + 3y}$$

So, the parametric curve will be some or all of the graph of this square root function.

Step 2

At this point let's get our first guess as to the limits on x and y. As noted in previous problems what we're really finding here is the largest possible ranges for x and y. In later steps we'll determine if this the actual set of limits on x and y or if we have smaller ranges.

We can use our knowledge of cosine to determine the limits on x and y as follows,

$$\begin{array}{ll} -1 \leq \cos\left(\frac{5}{2}t\right) \leq 1 & -1 \leq \cos\left(\frac{5}{2}t\right) \leq 1 \\ 3 \leq 4 + \cos\left(\frac{5}{2}t\right) \leq 5 & -\frac{1}{3} \leq \frac{1}{3}\cos\left(\frac{5}{2}t\right) \leq \frac{1}{3} \\ \sqrt{3} \leq \sqrt{4 + \cos\left(\frac{5}{2}t\right)} \leq \sqrt{5} & \frac{2}{3} \leq 1 + \frac{1}{3}\cos\left(\frac{5}{2}t\right) \leq \frac{4}{3} \\ \sqrt{3} \leq x \leq \sqrt{5} & \frac{2}{3} \leq y \leq \frac{4}{3} \end{array}$$

Remember that all we need to do is start with the cosine and then build up the equation for x and y by first multiplying the trig function by any coefficient, if present, and then adding/subtracting any numbers that might be present. We now have the largest possible set of limits for x and y.

Now, at this point we need to be a little careful. As noted above what we've actually found here are the largest possible ranges for the limits on x and y. This set of inequalities for the limits on x and y assume that the parametric curve will be fully traced out at least once for the range of t's we were given in the problem statement. It is always possible that the parametric curve will not trace out a full trace in the


given range of t 's. In a later step we'll determine if the parametric curve does trace out a full trace and hence determine the actual limits on x and y .

Remember that when we talk about the parametric curve getting fully traced out this doesn't, in general, mean the full square root graph we found in Step 1 gets traced out by the parametric equation. All "fully traced out" means, in general, is that whatever portion of the square root graph that is described by the set of parametric curves will be completely traced out.

In fact, for this problem, we can see that the square root from Step 1 will not get fully traced out by the particle regardless of any range of t 's. The largest possible portion of the square root graph that can be traced out by the particle is the portion that lies in the range of x and y given above. In a later step we'll determine if the largest possible portion of the square root graph does get traced out or if the particle only traces out part of it.

Step 3

Let's next get the direction of motion for the parametric curve. For this analysis it might be useful to have a quick sketch of the largest possible parametric curve. So, here is a quick sketch of that.



The dashed line is the graph of the full square root from Step 1 and the solid line is the portion that falls into our largest possible range of x and y we found in Step 2. As an aside here note that the two ranges are complimentary. In other words, if we sketch the graph only for the range of x we automatically get the range for y . Likewise, if we sketch the graph only for the range of y we automatically get the range for x . This is a good check for your graph. The x and y ranges should always match up!

Before moving on let's address the fact that it doesn't look like square root graphs that most of us are used to seeing. Keep in mind that the typical square root function that we're used to working at is in the form $y = \sqrt{x}$. Our equation for this problem however is in the form $x = \sqrt{y}$. If you think about it the graph of $x = \sqrt{y}$ is nothing more than the portion of the graph of $y = x^2$ corresponding to $x \geq 0$ (recall square roots only return positive or zero values!). Of course the function for this problem is not $x = \sqrt{y}$ but it is similar enough that the ideas discussed here are still valid just for a slightly different function.

Okay, let's get back to the problem.

This problem is going to be a lot like the previous problem in terms of direction of motion. First note that if we start at the lower left hand point we need to require that $\cos\left(\frac{5}{2}t\right) = -1$ since that is the only way for both x and y to have their minimal values (which puts us at the lower left-hand point)! It also doesn't matter what value of t we use at this point. All that matters is that we are at the lower left hand point.

If we now increase t (from whatever "starting" value we had) we know that cosine will need to increase from $\cos\left(\frac{5}{2}t\right) = -1$ until it reaches a value of $\cos\left(\frac{5}{2}t\right) = 1$. By looking at the parametric equations we can see that this will also force both x and y to increase until it reaches the upper right-hand point.

Now, the graph won't just stop here. Once cosine reaches a value of $\cos\left(\frac{5}{2}t\right) = 1$ we know that continuing to increase t will now cause cosine to decrease it reaches a value of $\cos\left(\frac{5}{2}t\right) = -1$. This in turn forces both x and y to decrease until it once again reaches the lower left-hand point.

In other words, if we don't put any restrictions on t a particle on this parametric curve will simply oscillate left and right along the portion of the square root sketched out above. In this case however we do have a range of t 's so we'll need to determine a range of t 's for one trace to fully know the direction of motion information of the particle on this path and we'll do that in the next step. With a restriction on the range of t 's it is possible that the particle won't make a full trace or it might retrace some or all of the curve so we can't say anything definite about the direction of motion for the particle over the full range of t 's until the next step when we determine a range of t 's for one full trace of the curve.

Step 4

Now we need to determine a range of t 's for one full trace of the parametric curve. It is important for this step to remember that one full trace of the parametric curve means that no portion of the parametric curve can be retraced.

Note that one full trace does not mean that we get back to the "starting" point. When we dealt with an ellipse in a previous problem that was one trace because we did not need to retrace any portion of the ellipse to get back to the starting point. However, as we saw in the previous step that for our square root here we would have to retrace the full curve to get back to the starting point.

So, one full trace of the parametric curve means we move from the right end point to the left end point only or visa-versa and move from the left end point to the right end point. Which direction we move doesn't really matter here so let's get a range of t 's that take us from the left end point to the right end point.

In order to be at the left end point we need to require that $\cos\left(\frac{5}{2}t\right) = -1$ which occurs if

$\frac{5}{2}t = \dots, -3\pi, -\pi, \pi, 3\pi, \dots$. Note as well that unlike the previous problems, which had both sine and cosine, this set of parametric equations has only cosine and so all we need to do here is look at this.

Also, in order to be at the right end point we need to require that $\cos\left(\frac{5}{2}t\right) = 1$ which occurs if

$\frac{5}{2}t = \dots, -4\pi, -2\pi, 0, 2\pi, 4\pi, \dots$

So, if we want to move from the left to right all we need to do is choose one from the list of t 's corresponding to the left end point and then first t that comes that from the list corresponding to the right end point and we'll have a range of t 's for one trace. To move from the right to left we just go the opposite direction, *i.e.* choose a t from the right end point list and then take the first t after that from the left end point list.

So, for this problem, since we said we were going to move from left to right, we'll use $\frac{5}{2}t = \pi$, which corresponds to $t = \frac{2}{5}\pi$, for the left end point. That in turn means that we'll need to use $\frac{5}{2}t = 2\pi$, which corresponds to $t = \frac{4}{5}\pi$, for the right end point. That means the range of t 's for one trace is,

$$\frac{2}{5}\pi \leq t \leq \frac{4}{5}\pi$$

This is only one possible answer here. Any range of t 's with a "net" range of $\frac{4}{5}\pi - (\frac{2}{5}\pi) = \frac{2}{5}\pi$ t 's, with the endpoints of the t range corresponding to start/end points of the parametric curve, will work. So, for example, any of the following ranges of t 's would also work.

$$-\frac{2}{5}\pi \leq t \leq 0 \qquad 0 \leq t \leq \frac{2}{5}\pi \qquad \frac{4}{5}\pi \leq t \leq \frac{6}{5}\pi$$

The direction of motion for each may be different range of t 's of course. Some will trace out the curve moving from left to right while others will trace out the curve moving from right to left. Because the problem did not specify a particular direction any would work.

Note as well that the range $\frac{2}{5}\pi \leq t \leq \frac{4}{5}\pi$ falls completely inside the given range of t 's specified in the problem and so we know that the particle will trace out the curve more than once over the full range of t 's. Determining just how many times it traces over the curve will be determined in the next step.

Step 5


Now that we have a range of t 's for one full trace of the parametric curve we can determine the number of traces the particle makes.

This is a really easy step. We know the total time the particle was traveling and we know how long it takes for a single trace. Therefore,

$$\text{Number Traces} = \frac{\text{Total Time Traveled}}{\text{Time for One Trace}} = \frac{2\pi - (-48\pi)}{\frac{4}{5}\pi - (\frac{2}{5}\pi)} = \frac{50\pi}{\frac{2}{5}\pi} = 125 \text{ traces}$$

Step 6

Finally, here is a sketch of the parametric curve for this set of parametric equations.



For this sketch we indicated the direction of motion by putting arrow heads going both directions in places on the curve. We also included a set of t 's for a couple of points to illustrate where the particle is at while tracing out of the curve as well as coordinates for the end points since they aren't "nice" points.. The dashed line is the continuation of the square root from Step 1 to illustrate that our parametric curve is only a part of the square root.

Here is also the formal answers for all the rest of the information that problem asked for.

Range of x :	$\sqrt{3} \leq x \leq \sqrt{5}$
Range of y :	$\frac{2}{3} \leq y \leq \frac{4}{3}$
Range of t for one trace :	$\frac{2}{5}\pi \leq t \leq \frac{4}{5}\pi$
Total number of traces :	125

10. The path of a particle is given by the following set of parametric equations. Completely describe the path of the particle. To completely describe the path of the particle you will need to provide the following information.

- (i) A sketch of the parametric curve (including direction of motion) based on the equation you get by eliminating the parameter.
- (ii) Limits on x and y .
- (iii) A range of t 's for a single trace of the parametric curve.
- (iv) The number of traces of the curve the particle makes if an overall range of t 's is provided in the problem.

$$x = 2e^t \quad y = \cos(1 + e^{3t}) \quad 0 \leq t \leq \frac{3}{4}$$

Step 1

There's a lot of information we'll need to find to fully answer this problem. However, for most of it we can follow the same basic ordering of steps we used for the first few problems in this section. We will need however to do a little extra work along the way.

Also, because most of the work here is similar to the work we did in Problems 4 – 6 of this section we won't be putting in as much explanation to a lot of the work we're doing here. So, if you need some explanation for some of the work you should go back to those problems and check the corresponding steps.

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations let's first notice that we can solve the x equation for the exponential function as follows,

$$e^t = \frac{1}{2}x$$

Now, just recall that $e^{3t} = (e^t)^3$ and so we can plug the above equation into the exponential in the y equation to get,

$$y = \cos\left(1 + e^{3t}\right) = \cos\left(1 + (e^t)^3\right) = \cos\left(1 + \left(\frac{1}{2}x\right)^3\right) = \cos\left(1 + \frac{1}{8}x^3\right)$$

So, the parametric curve will be some or all of the graph of this cosine function.

Step 2

At this point let's work on the limits for x and y . In this case, unlike most of the previous problems, things will work a little differently.

Let's start by noting that unlike sine and cosine functions we know e^t is always an increasing function (you can do some quick Calculus I work to verify this right?).

Why do we care about this? Well first the x equation is just a constant times e^t and we are given a range of t 's for the problem. Next, the fact that e^t is an increasing function means that the x equation, $x = 2e^t$, is also an increasing function (because the 2 is positive). Therefore, the smallest value of x will occur at the smallest value of t in the range of t 's. Likewise, the largest value of x will occur at the largest value of t in the range of t 's.

Therefore, the range of x for our parametric curve is,

$$2e^0 \leq x \leq 2e^{\frac{3}{4}} \quad \Rightarrow \quad 2 \leq x \leq 2e^{\frac{3}{4}}$$


Unlike the previous problems where we usually needed to do a little more verification work we know at this point that this is the range of x 's.

For the range of y 's we will need to do a little work to get the correct range of y 's but it won't be as much extra work as in previous problems and we can do it all in this step. First let's notice that because the y equation is in the form of $y = \cos(\dots)$. The argument of the cosine doesn't matter for the first part of the work and so wasn't included here.

From the behavior of cosine we then know that the largest possible range of y would then be,
$$-1 \leq y \leq 1$$

Now, depending on just what values the argument of the cosine in the y equation takes over the give range of t 's we may or may not cover this full range of values. We could do some work analyzing the argument of the cosine to figure that out if it does cover this full range. However, there is a really easy way to figure that if the full range is covered in this case.

Let's just sketch the graph and see what we get. Here is a quick sketch of the graph.



Given the “messy” nature of the argument of the cosine it’s probably best to use some form of computational aid to get the graph. The dotted portion of the graph is full graph of the function on $-3 \leq x \leq 5$ without regards to the actual restriction on t . The solid portion of the graph is the portion that corresponds to the range of t 's we were given in the problem.

From this graph we can see that the range of y 's is in fact $-1 \leq y \leq 1$.

Before proceeding with the direction of motion let’s note that we could also have just graphed the curve in many of the previous problems to determine if the work in this step was the actual range or not. We didn’t do that because we could determine if these ranges were correct or not when we did the direction of motion and range of t 's for one trace analysis (which we had to do anyway) and so didn’t need to bother with a graph in this step for those problems.

Step 3

We now need to do the direction of motion for this curve but note that we actually found the direction of motion in the previous step.


As noted in the previous step we know that $x = 2e^t$ is an increasing function and so the x 's must be increasing as t increases. Therefore, the equation must be moving from left to right as the curve is traced out over the given range of t 's.

Also note that unlike the previous problems we know that no portion of the graph will be retraced. Again, we know the x equation is an increasing equation. If the curve were to retrace any portion we can see that the only way to do that would be to move back from right to left which would require x to decrease and that can’t happen.

This means that we now know as well that the graph will trace out exactly once for the given range of t 's, which in turn tells us that the given range of t 's is also the range of t 's for a single trace.

Step 4

Now that we have all the needed information we can do a formal sketch of the graph.



As with the graph above the dotted portion of the graph is full graph of the function on $-3 \leq t \leq 5$ without regards to the actual restriction on t . The solid portion of the graph is the portion that corresponds to the range of t 's we were given in the problem. We also included the t value and coordinates of each end point for clarity although these are often not required for many problems.

Here is also the formal answers for all the rest of the information that problem asked for.

Range of x :	$2 \leq x \leq 2e^{\frac{3}{4}}$
Range of y :	$-1 \leq y \leq 1$
Range of t for one trace :	$0 \leq t \leq \frac{3}{4}$
Total number of traces :	1

11. The path of a particle is given by the following set of parametric equations. Completely describe the path of the particle. To completely describe the path of the particle you will need to provide the following information.

(i) A sketch of the parametric curve (including direction of motion) based on the equation you get by eliminating the parameter.

(ii) Limits on x and y .

(iii) A range of t 's for a single trace of the parametric curve.

(iv) The number of traces of the curve the particle makes if an overall range of t 's is provided in the problem.

$$x = \frac{1}{2}e^{-3t} \quad y = e^{-6t} + 2e^{-3t} - 8$$

Step 1

There's a lot of information we'll need to find to fully answer this problem. However, for most of it we can follow the same basic ordering of steps we used for the first few problems in this section. We will need however to do a little extra work along the way.

Also, because most of the work here is similar to the work we did in Problems 4 – 6 of this section we won't be putting in as much explanation to a lot of the work we're doing here. So, if you need some explanation for some of the work you should go back to those problems and check the corresponding steps.

First, we'll eliminate the parameter from this set of parametric equations. For this particular set of parametric equations let's first notice that we can solve the x equation for the exponential function as follows,

$$e^{-3t} = 2x$$

Now, just recall that $e^{-6t} = (e^{-3t})^2$ and so we can plug the above equation into the exponential in the y equation to get,

$$y = e^{-6t} + 2e^{-3t} - 8 = (e^{-3t})^2 + 2e^{-3t} - 8 = (2x)^2 + 2(2x) - 8 = 4x^2 + 4x - 8$$

So, the parametric curve will be some or all of the graph of this quadratic function.

Step 2

At this point let's work on the limits for x and y . In this case, unlike most of the previous problems, things will work a little differently.

Let's start by noting that unlike sine and cosine functions we know e^{-3t} is always a decreasing function as t increases (you can do some quick Calculus I work to verify this right?).

Why do we care about this? Well first the x equation is just a constant times e^{-3t} and so the fact that e^{-3t} is a decreasing function means that the x equation, $x = \frac{1}{2}e^{-3t}$, is also a decreasing function (because the coefficient is positive).

Next, we aren't given a range of t 's for this problem and so we can assume the largest possible range of t 's. Therefore, we are safe in assuming a range of $-\infty < t < \infty$ for the t 's.

Now, as we've already noted we know that the x equation is decreasing and so the largest value of x will occur at the left "end point" of the range. Likewise, the smallest value of x will occur at the right "end point" of the range. For this problem both "end points" of our range are in fact infinities so we can't just plug in as we did in the previous problem. We can however take the following two limits.

$$\lim_{t \rightarrow -\infty} \left(\frac{1}{2} e^{-3t} \right) = \infty$$

$$\lim_{t \rightarrow \infty} \left(\frac{1}{2} e^{-3t} \right) = 0$$


From this we can see that as we approach the left end point of the t range the value of x is going to infinity and as we approach the right end point of the t range the x value is going to zero. Note however that x can never actually be zero because x is still defined in terms of an exponential function (which can't be zero). All the limit is telling us is that as we let $t \rightarrow \infty$ we will get $x \rightarrow 0$.

The range of x for our parametric curve is therefore,

$$0 < x < \infty$$

Again, be careful with the inequalities here! We know that x can be neither zero nor infinity so we must use strict inequalities for this range. This is something that we always need to be on the lookout for with variable ranges of parametric equations. Depending on the parametric equations sometimes the end points of the ranges will be strict inequalities (as with this problem) and for others they include the end points (as with the previous problems).

For the range of y 's we will need to do a little work to get the correct range of y 's but it won't be as much extra work as in previous problems and we can do it all in this step. Let's just sketch the graph and see what we get. Here is a quick sketch of the graph.



The dotted portion of the graph is full graph of the function on $-2 \leq t \leq 2$ without regards to the actual restriction on x . The solid portion of the graph is the portion of the graph that corresponds to the restriction on x that we found earlier in this step.

Note the "open dot" on the y -axis for the left end of the graph. This needs to be here to acknowledge that $x \neq 0$. We can also see that the y value at this point is $y = -8$ and again we can see that for the parametric curve we have $y \neq -8$.

Also, keep in mind that we know that $x \rightarrow \infty$ and so we also know that the y portion of the graph must also continue up to infinity to match the x behavior.

So, from this quick analysis of the graph we can see that the y range for the parametric curve must be,

$$-8 < y < \infty$$

Step 3

We now need to do the direction of motion for this curve but note that we actually found the direction of motion in the previous step.


As noted in the previous step we know that $x = \frac{1}{2}e^{-3t}$ is a decreasing function and so the x 's must be decreasing as t increases. Therefore, the equation must be moving from right to left as the curve is traced out.

Also note that unlike most of the previous problems we know that no portion of the graph will be retraced. Again, we know the x equation is a decreasing equation. If the curve were to retrace any portion we can see that the only way to do that would be to move back from left to right which would require x to increase and that can't happen.

This means that we now know as well that the graph will trace out exactly once.

Step 4

Now that we have all the needed information we can do a formal sketch of the graph.



As with the graph above the dotted portion of the graph is full graph of the function on $-2 \leq t \leq 2$ and the solid portion of the graph is the portion that corresponds to the restrictions x and y we found in Step 2.

Here is also the formal answers for all the rest of the information that problem asked for.

$$\text{Range of } x : \quad 0 \leq x < \infty$$

$$\text{Range of } y : \quad -8 \leq y < \infty$$

$$\text{Range of } t \text{ for one trace} : \quad -\infty < t < \infty$$

$$\text{Total number of traces} : \quad 1$$

12. Write down a set of parametric equations for the following equation.

$$y = 3x^2 - \ln(4x + 2)$$

Solution

There really isn't a lot to this problem. All we need to do is use the "formulas" right at the end of the notes for this section.

A set of parametric equations for the equation above are,

$$\begin{aligned}x &= t \\ y &= 3t^2 - \ln(4t + 2)\end{aligned}$$

13. Write down a set of parametric equations for the following equation.

$$x^2 + y^2 = 36$$

The parametric curve resulting from the parametric equations should be at $(6, 0)$ when $t = 0$ and the curve should have a counter clockwise rotation.

Solution

If we don't worry about the "starting" point (*i.e.* where the curve is at when $t = 0$) and we don't worry about the direction of motion we know from the notes that the following set of parametric equations will trace out a circle of radius 6 centered at the origin.

$$\begin{aligned}x &= 6 \cos(t) \\ y &= 6 \sin(t)\end{aligned}$$

All we need to do is verify if the extra requirements are met or not.

First, we can clearly see with a quick evaluation that when $t = 0$ we are at the point $(6, 0)$ as we need to be.

Next, we can either use our knowledge from the examples worked in the notes for this section or an analysis similar to some of the earlier problems in this section to verify that circles in this form will always trace out in a counter clockwise rotation.

In other words, the set of parametric equations give above is a set of parametric equations which will trace out the given circle with the given restrictions. So, formally the answer for this problem is,

$$\boxed{\begin{aligned}x &= 6 \cos(t) \\ y &= 6 \sin(t)\end{aligned}}$$

We'll leave this problem with a final note about the answer here. This is possibly the "simplest" answer we could give but it is completely possible that you may have come up with a different answer to this problem. There are almost always lots of different possible sets of parametric equations that will trace out a particular parametric curve according to some particular set of restrictions.

14. Write down a set of parametric equations for the following equation.

$$\frac{x^2}{4} + \frac{y^2}{49} = 1$$

The parametric curve resulting from the parametric equations should be at $(0, -7)$ when $t = 0$ and the curve should have a clockwise rotation.

Solution

If we don't worry about the "starting" point (*i.e.* where the curve is at when $t = 0$) and we don't worry about the direction of motion we know from the notes that the following set of parametric equations will trace out the ellipse given by the equation above.

$$\begin{aligned}x &= 2 \cos(t) \\y &= 7 \sin(t)\end{aligned}$$

The problem with this set of parametric equations is that when $t = 0$ we are at the point $(7, 0)$ which is not the point we are supposed to be at. Also, from our knowledge of the examples worked in the notes for this section or an analysis similar to some of the earlier problems in this section we can see that the parametric curve traced out by this set of equations will trace out in a counter clockwise rotation – again not what we need.

So, we need to come up with a different set of parametric equations that meets the requirements.

The first thing to acknowledge is that using sine and cosine will always be the easiest way to get a set of parametric equations for an ellipse. However, there is no reason at all to always use cosine for the x equation and sine for the y equation.

Knowing that we need $x = 0$ and $y = -7$ when $t = 0$ and using the fact that we know that $\sin(0) = 0$ and $\cos(0) = 1$ the following set of parametric equations will "start" at the correct point when $t = 0$.

$$\begin{aligned}x &= -2 \sin(t) \\y &= -7 \cos(t)\end{aligned}$$

All we need to do now is check if this will trace out the ellipse in a clockwise direction.

If we start at $t = 0$ and increase t until we reach $t = \frac{\pi}{2}$ we know that sine will increase from 0 to 1. This will in turn mean that x must decrease (don't forget the minus sign on the x equation) from 0 to -2.

Likewise, increasing t from $t = 0$ to $t = \frac{\pi}{2}$ we know that cosine will decrease from 1 to 0. This in turn means that y will increase (don't forget the minus sign on the y equation!) from -7 to 0.

The only way for both of these things to happen at the same time is for the curve to start at $(0, -7)$ when $t = 0$ and trace along the ellipse in a clockwise direction until we reach the point $(-2, 0)$ when $t = \frac{\pi}{2}$.

We could continue in this fashion further increasing t until it reaches $t = 2\pi$ (which will put us back at the "starting" point) and convince ourselves that the ellipse will continue to trace out in a clockwise direction.

Therefore, one possible set of parametric equations that we could use is,

$$\begin{array}{l} x = -2 \sin(t) \\ y = -7 \cos(t) \end{array}$$

We'll leave this problem with a final note about the answer here. This is possibly the "simplest" answer we could give but it is completely possible that you may have come up with a different answer to this problem. There are almost always lots of different possible sets of parametric equations that will trace out a particular parametric curve according to some particular set of restrictions.
